

TALPRID®



FIRST AID
HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE
IF SWALLOWED:

- Call a poison control center, doctor, or 1-877-864-6384 immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

IF ON SKIN:

- Wash with plenty of soap and water.

TREATMENT FOR PET FEEDING
 If an animal eats bait, call veterinarian or 1-877-454-6384 at once.

NOTE TO VETERINARIAN OR VETERINARIAN
 Contains the nerve poison Bromethalin. This is not an anticoagulant type rodenticide. For humans or animals regarding bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), full absorption by other animals or aquatic biota. Significant symptoms, if present, would be the result of oral exposure and should be treated accordingly through administration of an emetic, diazepam and corticosteroid.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children to the product or use it for any purpose other than to control moles. To help prevent accidents:

1. Store product not in use in locations out of the reach of children and pets.
2. Follow all application directions and USE RESTRICTIONS on this label. Apply bait only in underground runways of moles. Keep pets out of treated areas. Do NOT use the product above ground.
3. Dispose of product container and unused or spoiled product as specified on label.

USE RESTRICTIONS: This product may only be applied at the use sites and by the application methods indicated on this label. This product may only be used in lawns, parks, around homes, golf courses, commercial gardens, lawns and other non-crop grassy areas to control eastern moles (*Scalopus aquaticus*), star-nosed mole (*Condalia cristata*), and Skink-footed mole (*Scalopus aquaticus*). Do not apply directly into mole tunnel systems. This bait must be applied underground in either main underground runways or in subsurface feeding tunnels. This bait may not be used above ground, near, over, or near bait is applied.

SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of subsurface runways in the turf or by a series of conical mounds of earth pushed up from deep burrows. Subsurface runways are recognized by raised ridges of vegetation and soil caused by mole digging for food just below the soil surface. These runways are especially noticeable when the grass above them turns yellow. Conical mounds may be indicative of the location of deeper main underground runways.

APPLICATION DIRECTIONS:
 Remove soil and/or remove bait according to use directions below.

SUBSURFACE RUNWAYS: TALPRID® has been specially designed to mimic the natural food sources of the mole. To achieve maximum effect, TALPRID® should only be placed in areas used habitually by moles for feeding.

ESTABLISHING INITIAL ACTIVITY: Use your finger, small wooden dowel or a non-metallic object to make a hole in the top of subsurface runways. Be careful not to crush runways. Mark opened runways and resist them 48 to 72 hours later. Runways that have had holes inserted within 72 hours should be baited.

BAITING ACTIVE RUNWAYS: Make a hole in the top of the runway with a rod slightly larger than the diameter of the bait. Drop one worm-shaped bait into the runway using the rod to push any exposed bait completely into the runway. Carefully reseal the runway with a small amount of dirt to exclude light. Repeat bait application every 5 to 10 feet of each active subsurface runway.

DETERMINING BAITING RESULTS: Five (5) to 7 days after bait application, check for mole activity using the same procedure for determining initial activity. Return 48 to 72 hours later and retreat all active runs.

MAIN UNDERGROUND TUNNELS AND CONICAL MOUNDS: Mole's construct deeper tunnels that are not visible from the surface. These deep tunnels systems are often marked by a series of conical mounds (piles of dirt) 4 to 12 inches high formed when dirt is pushed to the surface when moles construct these tunnels. To locate deep tunnels, push a probe, such as a broom handle, into the soil between mounds. When this probe enters a tunnel, the resistance on the probe will decrease rapidly. Mark opened tunnels and resist them 48 to 72 hours later. Runways that have had holes inserted within 72 hours should be baited.

To bait deep tunnels, drop one TALPRID® worm through the probe hole, using the probe to push the bait completely into the tunnel. Cover the hole with a piece of soil, a small rock or a piece of newspaper to allow for visual inspection. Repeat bait application every 5 to 10 feet of each active deep tunnel and within 5 feet of each active conical mound. Check for the effects of treatment 5 to 7 days after bait application using the same procedure for determining tunnel location. Return 48 to 72 hours later and retreat all active tunnel systems.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin, or on clothing.

All handlers (including applicators) must wear shoes, socks, and gloves. Any person who inhales, consumes or uses bait before application of this product must wear eye protection.

User Safety Requirements
 Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washable, use detergent and hot water. Stop and wash PPE immediately after handling this product. Wash the outside of boots before removing. As soon as possible, wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet and changing into clean clothing.

Prolonged or frequent repeat skin contact may cause allergic reactions in some individuals.

Environmental Hazards
 The product is extremely toxic to mammals and birds. Dogs and other predatory and scavenging mammals, rodents and birds should be kept away from animals that have eaten this bait. Do not apply directly to water.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.
PESTICIDE DISPOSAL: Pests resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.
CONTAINER HANDLING: Handle with care. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container by placing in trash, or in an approved waste disposal facility or incineration.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling goes contrary to label restrictions.

11191800-015
 Manufactured by
Bell
LABORATORIES, INC.
 3689 Krummer Blvd.
 Madison, WI 53704-USA
 www.belllab.com
 MADE IN USA


* Not for use against the star-nosed mole in North Carolina

KILLS MOLES



20 WORM SHAPED BAITS

TALPRID®



KILLS MOLES
MIMICS A MOLE'S NATURAL FOOD SOURCE, THE EARTHWORM
READY TO USE

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See back panel for important information.
 See back panel for important information.
 See back panel for important information.

NET WT:
3.5 OZ (100 g)
 Contains 20 worm shaped baits
 (TWO Individually Sealed Trays of 10)

KILLS MOLES



2185

FIRST AID	
HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE	
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.
IF ON SKIN:	<ul style="list-style-type: none"> Wash with plenty of soap and water.

TREATMENT FOR PET POISONING	
If an animal eats bait, call veterinarian or 1-877-854-2494 at once.	
NOTE TO PHYSICIAN OR VETERINARIAN	
Contains the nerve poison Bromethalin. This is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.	

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children to this product or use it for any purpose other than to control moles. To help prevent accidents:

1. Store product not in use in locations out of the reach of children and pets.
2. Follow all application directions and USE RESTRICTIONS on this label. Apply bait only in underground runways of moles. Keep pets out of treated areas. Do NOT use this product above ground.
3. Dispose of product container and unused or spoiled product as specified on label.

USE RESTRICTIONS: This product may only be applied at the use sites and by the application methods indicated on this label. This product may only be used in lawns, parks, around homes, golf courses, ornamental gardens, nurseries and other non-crop grassy areas to control eastern moles (*Scalopus aquaticus*), star-nosed moles* (*Corydora cristata* parva) or *Scapanus* spp. Bait must be applied directly into moles' burrow systems. This bait must be applied underground in either main underground runways or in subsurface feeding tunnels. This bait may not be used above ground. Wear gloves when bait is handled.

SELECTION OF TREATMENT AREAS: The presence of moles may be indicated by a network of subsurface runways in the turf or by a series of conical mounds of earth pushed up from deep burrows. Subsurface runways are indicated by raised ridges of vegetation and soil caused by moles foraging for food just below the soil surface. These runways are especially noticeable when the grass above them turns yellow. Conical mounds may be indicative of the location of deeper main underground runways.

APPLICATION DIRECTIONS:
Remove and place worm shaped bait according to use directions below.

SUBSURFACE RUNWAYS: TALPIRID® has been specially designed to mimic the natural food sources of the mole. To achieve maximum effect, TALPIRID® should only be placed in areas used habitually by moles for feeding.

751MO-8 * Not for use against the star-nosed mole in North Carolina

TALPIRID®

KILLS MOLES

This Product May Only be Used to Control Moles* in Manual, Below-Ground Applications.
EFFECTIVE AGAINST MOST COMMON MOLE SPECIES
MIMICS A MOLE'S NATURAL FOOD SOURCE, THE EARTHWORM

READY TO USE
Scientifically designed and tested to mimic a mole's natural food source. Moles may consume a lethal dose in a single feeding, but it may take two or more days from the time of bait consumption for moles to die.
ACTIVE INGREDIENT: Bromethalin (CAS #63333-35-7) 0.025%
OTHER INGREDIENTS: 99.975%
TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling are contrary to label instructions.

See side panels for First Aid and additional precautionary statements.

Contains 10 Worm Shaped Baits
Manufactured by: **Bell Laboratories, Inc.**
3600 Kinsman Blvd.
Madison, WI 53704 U.S.A.
www.belllabs.com

NET WEIGHT:
1.76 oz (50 g)

EPA REG. NO. 12455-101
EPA EST. NOS. 12455-WI-K
12455-WI-2P, 12455-WI-3M
Superscript is first letter of the lot number.

INDIVIDUAL SALE IS PROHIBITED BY LAW

DO NOT EXPOSE TO EXCESSIVE HEAT STORE BELOW 100° F STORE FLAT - THIS SIDE UP

DIRECTIONS FOR USE (Continued from other panel)

ESTABLISHING INITIAL ACTIVITY: Use your finger, small wooden dowel or a narrow rod to puncture a hole in the top of subsurface runways. Be careful not to crush runways. Mark opened runways and revisit them 48 to 72 hours later. Runways that have had holes resealed within 72 hours should be baited.

BAITING ACTIVE RUNWAYS: Make a hole in the top of the runway with a rod slightly larger than the diameter of the bait. Drop one worm into the runway, using the rod to push any exposed bait completely into the runway. Carefully reseat the runway with a small amount of dirt to exclude light. Repeat bait application every 5 to 10 feet of each active subsurface runway.

DETERMINING BAITING RESULTS: Five (5) to 7 days after bait application, check for mole activity using the same procedure for determining initial activity. Return 48 to 72 hours later and retreat all active runs.

MAIN UNDERGROUND TUNNELS AND CONICAL MOUNDS: Moles construct deeper tunnels that are not visible from the surface. These deep tunnels systems are often marked by a series of conical mounds (piles of dirt) 4 to 12 inches high formed when dirt is pushed to the surface when moles construct these tunnels. To locate deep tunnels, push a probe, such as a broom handle, into the soil between mounds. When the probe enters a tunnel, the resistance on the probe will decrease rapidly. Mark opened tunnels and revisit them 48 to 72 hours later. Runways that have had holes resealed within 72 hours should be baited.

To bait deep tunnels, drop one TALPIRID® worm through the probe hole, using the probe to push the bait completely into the tunnel. Cover the hole with a piece of sod, a small rock or a piece of newspaper to allow for visual inspection. Repeat bait application every 5 to 10 feet of each active deep tunnel and within 5 feet of each active conical mound. Check for the effects of treatment 5 to 7 days after bait application using the same procedure for determining tunnel location. Return 48 to 72 hours later and retreat all active tunnel systems.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin, or on clothing.

All handlers (including applicators) must wear shoes, socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

User Safety Requirements
Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

Prolonged or frequent repeat skin contact may cause allergic reactions in some individuals.
Environmental Hazards
This product is extremely toxic to mammals and birds. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

112018/05-15